



Faculty of Resource Science and Technology

**EFFECT OF REACTION TIME ON BASE HYDROLYSIS
OF METHYL BENZOATE**

ROSSIDAH SINJA ANAK DAVID DARI

(43864)

**Bachelor of Science with Honours
(Resource Chemistry)
2016**

EFFECT OF REACTIONN TIME ON BASE HYDROLYSIS OF METHYL BENZOATE

ROSSIDAH SINJA ANAK DAVID DARI (43864)

**This project paper is submitted in partial fulfillment of requirements the Degree of
Bachelor of Science with honors
(Program of Resource Chemistry)**

Faculty of Resource Science and Technology

UNIVERSITI MALAYSIA SARAWAK

2016

Acknowledgement

I would like to express my deepest gratitude to the many people that helped to bring this research project to success. I want to thank, first and foremost my supervisor, Miss Yusralina bt Yusof for providing me the opportunity to taking part in this Final Year Project. I am so grateful for her help, professionalism and valuable guidance throughout this project that I do not have enough words to give my sincere appreciation.

I would also like to thank all the staffs that involve in the analysis of data for this research project. Without their passionate participation and input, the analysis of data could not have successfully conducted.

I am indebted to my friends who support me in this research project through the entire year. I cannot find words to express my gratitude to them in making this research project to fruition. I consider it an honor to work with all of them during this project.

Last but not least, I must express my profound gratitude to my parents for providing me with unfailing support throughout my years of study and through the process of researching and writing this thesis. This thesis would not have been possible without them.

Thank you.

Grade: A

Please tick (✓)

Final Year Project Report ☒

Masters ☐

PhD ☐

DECLARATION OF ORIGINAL WORK

This declaration is made on the 17th May 2012.

Student's Declaration:

I Rossidah Sinja anak David Dari, 43864, Faculty of Resource Science and Technology hereby declare that the work entitled, Effect of Reaction Time on Base Hydrolysis of Methyl Benzoate is my original work. I have not copied from any other students' work or from any other sources except where due reference or acknowledgement is made explicitly in the text, nor has any part been written for me by another person.

17th May 2012

Date submitted

RS


Rossidah Sinja anak David Dari (43864)

Supervisor's Declaration:

I, Yusralina binti Yusof, hereby certify that the work entitled, Effect of Reaction Time on Acid Hydrolysis of Methyl Benzoate was prepared by the above name student, and was submitted to the "FACULTY" as a full fulfilment for the conferment of Bachelor of Science

with Honours (Resource Chemistry), and the aforementioned work, to the best of my knowledge, is the said student's work.

Received for examination by:


(Yusralina binti Yusof)

Date: 27 / 6 / 2016

I declare this Project/Thesis is classified as (Please tick (√)):

☐

CONFIDENTIAL (Contain confidential information under the Official Secret Act 1972)*

☐

RESTRICTED (Contains restricted information as specified by the organisation where research was done)*

☐

OPEN ACCESS


Validation of Project/Thesis

I therefore duly affirmed with free consent and willingness declared that this said Project/Thesis shall be placed officially in the Centre for Academic Information Services with the abide interest and rights as follows:


- This Project/Thesis is the sole legal property of Universiti Malaysia Sarawak (UNIMAS).
- The Centre for Academic Information Services has the lawful right to make copies for the purpose of academic and research only not for other purposes.
- The Centre for Academic Information Services has the lawful right to digitise the content to for the Local Content Database.
- The Centre for Academic Information Services has the lawful right to make copies of the Project/Thesis for the academic exchange between Higher Learning Institute.

- No dispute or any claim shall arise from the student himself / herself neither third party on this Project/Thesis once it becomes sole property of UNIMAS.
- This Project/Thesis or any material, data and information related to it shall not be distributed, published or disclosed to any party by the student except with UNIMAS permission.

Student's signature: _____


(17th May 2016)

Supervisor's signature: _____


(17th May 2016)

Current Address:

Rumah Sandah Ulu Machan

96700, Kanowit

Notes: * If the Project/Thesis is COFIDENTIAL or RESTRICTED, please attach together as annexure a letter from the organisation with the period and reasons of confidentiality and restriction.

Table of Content

Acknowledgement III

DECLARATION OF ORIGINAL WORKIV

Table of Content VII

List of Abbreviation..... X

List of Table.....XI

List of Figure XII

Abstract.....XIV

1.0 Introduction..... 1

1.1 Background..... 1

1.2 Problem statement.....3

1.3 Objective 3

2.0 Literature Review4

2.1 Ester..... 4

2.1.1 Methyl benzoate4

2.2 Benzoic acid..... 5

2.2.1. Application of benzoic acid.....6

2.3 Hydrolysis6

2.3.1 Base hydrolysis.....7

2.3.2 Base hydrolysis of carboxylic ester7

2.3.2 Acid hydrolysis..... 8

2.3.3 Solvent effects on hydrolysis.....8

3.0 Materials and Methods.....	10
3.1 Preparation of 2 M NaOH.....	10
3.3 Preparation of 2 M HCl.....	10
3.4 Base hydrolysis of methyl benzoate with solvent.....	10
3.5 Purification of benzoic acid	11
3.6 Preparation of sample for analysis	11
3.6.1 FTIR analysis.....	11
3.6.2 Melting point analysis	12
4.0 Result and Discussion.....	13
4.1 Effect off reaction time on base hydrolysis	13
4.1.1 Comparison between the effect of reaction time on methyl benzoate by using NaOH in the presence and absence of MeOH.....	17
4.1.2 Comparison between the effect of reaction time on base hydrolysis of methyl benzoate by using KOH in the presence and absence of MeOH.....	18
4.1.3 Comparison between the effect of reaction time on methyl benzoate by using NaOH and KOH in the presence of MeOH.....	19
4.1.4 Comparison between the effect of reaction time on methyl benzoate by using NaOH and KOH in the absence of MeOH	20
4.2 FTIR analysis of benzoic acid.....	21
4.3 NMR analysis of benzoic acid	22
4.3.1 ^1H NMR analysis.....	22
4.3.2 ^{13}C NMR analysis.....	23

4.4 Melting point analysis of benzoic acid24

5.0 Conclusion26

6.0 References.....27

7.0 Appendix.....28

EtOH	Ethanol
MeOH	Methanol
MeOTf	Methanetrioxofluoride
HOAc	Acetic Acid
PhBr	Bromobenzene

List of Abbreviation

C=O	Carbonyl
FTIR	Fourier Transform Infrared Spectroscopy
NMR	Nuclear Magnetic Resonance
KOH	Potassium Hydroxide
NaOH	Sodium Hydroxide
MeOH	Methanol
HCl	Hydrochloric Acid
KBr	Potassium Bromide

List of Table

Table 1: Effect of reaction time on base hydrolysis of methyl benzoate by using NaOH in the presence of MeOH	13
Table 3: Effect of reaction time on base hydrolysis of methyl benzoate by using KOH in the presence of MeOH	15
Table 4: Effect of reaction time on base hydrolysis of methyl benzoate by using KOH in the absence of MeOH	16
Table 5: ^1H NMR data	23
Table 6: ^{13}C NMR	24
Table 7: Melting point of benzoic acid	24
Table 8: Experimental data for hydrolysis of methyl benzoate to form benzoic acid	28
Table 9: Base hydrolysis of methyl benzoate by using NaOH in the presence of MeOH	28
Table 10: Base hydrolysis of methyl benzoate by using NaOH as catalyst in the absence of MeOH	29
Table 11: Base hydrolysis of methyl benzoate by using KOH as catalyst in the absence of MeOH	29
Table 12: Base hydrolysis of methyl benzoate by using KOH as catalyst in the presence of MeOH	30

List of Figure

Figure 1: Structure of methyl benzoate.....	1
Figure 2: Transesterification of methyl benzoate to propyl acetate.....	1
Figure 3: Structure of benzoic acid.....	2
Figure 4: Mechanism of chlorinating of toluene to give benzoic acid	2
Figure 5: Fischer's esterification	5
Figure 6: Reaction of methyl benzoate with ammonia.....	5
Figure 7: Oxidation of toluene with air	5
Figure 8: Preparation of benzoic acid from phthalic anhydride	6
Figure 9: <i>Equation of hydrolysis process</i>	7
Figure 10: Hydrolysis reaction of oils and fats.....	7
Figure 11: Mechanism of base hydrolysis of carboxylic ester	8
Figure 12: Mechanism of acid hydrolysis.....	8
Figure 13: Pathway of hydrolysis of benzoic acid from methyl benzoate	10
Figure 14: Effect of reaction time on base hydrolysis of methyl benzoate by using NaOH in the presence of MeOH	13
Figure 15: Effect of reaction time on base hydrolysis of methyl benzoate by using NaOH in the absence of MeOH	14
Figure 16: Effect of reaction time on base hydrolysis of methyl benzoate by using KOH in the presence of MeOH	15
Figure 17: Effect of reaction time on base hydrolysis of methyl benzoate by using KOH in the absence of MeOH	16
Figure 18: Comparison of percentage yield for base hydrolysis by using NaOH in the presence and absence of MeOH.....	17

Figure 19: Comparison of percentage yield for base hydrolysis by using KOH in the presence and absence of MeOH.....	18
Figure 20: Comparison of percentage yield for base hydrolysis by using NaOH and KOH in the presence of MeOH.....	19
Figure 21: Comparison of percentage yield for base hydrolysis by using NaOH and KOH in the presence of MeOH.....	20
Figure 22: FTIR analysis of benzoic acid.....	21
Figure 23: ^1H NMR analysis of benzoic acid	22
Figure 24: ^{13}C NMR analysis of benzoic acid.....	23

Abstract

Hydrolysis is a reaction with water in general where the initially one compound converts to two compounds. Hydrolysis of methyl benzoate produced benzoic acid. Previous study found that the hydrolysis of *Jatropha curcas* seed oil to give free fatty acid (FFA) increase with increasing time. Therefore, this study aims to determine the effect of reaction time on base hydrolysis of methyl benzoate by using NaOH and KOH as catalyst. The effect of reaction time for base hydrolysis of benzoic acid by using NaOH and KOH as catalyst has been compared. Different reaction times (1.0, 2.0, 5.0, 7.0, 13.0 and 18.0) were used to determine the optimum yield in the presence and absence of methanol in base hydrolysis. The result obtained shown that percentage yield increase with increasing time.

Key Words: Base hydrolysis, Time, Benzoic acid, NaOH, KOH

Abstrak

Hidrolisis adalah tindak balas dengan air secara umum di mana satu kompaun pada mulanya menjadi dua sebatian. Hidrolisis methyl benzoate menghasilkan asid benzoik. Kajian sebelum ini mendapati bahawa hidrolisis minyak biji *Jatropha curcas* untuk menghasilkan asid lemak bebas (FFA) meningkat dengan peningkatan masa. Oleh itu, kajian ini bertujuan untuk menentukan kesan masa tindak balas pada hidrolisis methyl benzoate dengan menggunakan NaOH dan KOH sebagai pemangkin. Kesan masa tindak balas untuk hidrolisis alkali asid benzoik dengan menggunakan NaOH dan KOH sebagai pemangkin telah dibandingkan. Masa tindak balas yang berbeza (1.0, 2.0, 5.0, 7.0, 13.0 dan 18.0) telah digunakan untuk menentukan hasil optimum dengan kehadiran dan ketiadaan metanol dalam hidrolisis alkali. Keputusan yang diperolehi menunjukkan bahawa peningkatan hasil peratusan meningkat dengan peningkatan masa.

Kata kunci: Hidrolisis alkali, Masa, asid benzoik, NaOH, KOH

1.0 Introduction

1.1 Background

Methyl benzoate (Figure 1) is an ester that derived from benzoic acid (Marumaya *et al.*, 2012). It is a compound that has an ester group attached to a benzene ring. According to Marumaya *et al.* (2012), methyl benzoate plays an important role in the intermediate production of dyes, flavoring compounds, and mold growth biomarkers. The study found that methyl benzoate is important in the production of higher ester through transesterification in which methyl benzoate reacts with high boiling point alcohol such as butanol and benzyl alcohol to give benzyl or butyl benzoate and methanol as a product (Figure 2).




Figure 1: Structure of methyl benzoate




Figure 2: Transesterification of methyl benzoate to propyl acetate

Benzoic acid (Figure 3) is a compound that has a carboxylic group attached to a benzene ring. According to Bahl & Bahl (2006), benzoic acid is a white solid which is soluble in pure solvent with a melting point of 122°C and a boiling point of 249°C. The authors stated that benzoic acid is slightly soluble in cold water but dissolves completely in hot water. In addition, benzoic acid is important in various applications such as in medicines, food preservative and in dye industry. It is also used in cosmetic products (Nair, 2001). Benzoic acid can be prepared through various preparations such as oxidation of toluene, chlorinating of toluene followed by hydrolysis to give benzoic acid (Figure 4) and through reaction of phthalate anhydride in the presence of metal phthalate as catalyst (Bahl & Bahl,

2006). Benzoic acid should be handled carefully as the vapours could lead to the coughing and sneezing due to its pungent odor (Bahl & Bahl, 2006).




Figure 3: Structure of benzoic acid




Figure 4: Mechanism of chlorinating of toluene to give benzoic acid

Benzoic acid was synthesized from methyl benzoate by base hydrolysis in the present study using NaOH and KOH as catalyst in this study. According to Bruice (2006), hydrolysis reaction is a reaction with water where the initially one compound converts to two compounds. The study stated that ester was hydrolysed slowly because water is a poor nucleophile while the ester is a poor leaving group. However, the rate of reaction can be increased in the presence of acid or base as the catalyst. The one that used acid is known as acid hydrolysis while the one that used base is known as base hydrolysis. There are various organic compounds other than esters that undergo hydrolysis reaction such as amides, alkyl halides and epoxides (Alhamary *et al.*, 2014).

1.2 Problem statement

The study done by Al-Bahri *et al.* (2009) and Salimon *et al.* (2011) found that the percentage yield is increased with increasing time for the hydrolysis of ester. According to Salimon *et al.* (2011), in the hydrolysis of *Jatropha curcas* seed oil to produce free fatty acid (FFA) three different time have been used (1.5, 2.0 and 2.5 hours). The percentage of FFA is increase with time. Among these reaction times, 2.0 hours showed the highest percentage yield of FFA. Therefore, this study aimed to determine the optimum time to give the maximum yield through the process of base hydrolysis. In this study, methyl benzoate was hydrolyzed in the presence of NaOH and KOH as catalyst. The base hydrolysis was conducted at different reaction times which are 1.0, 2.0, 5.0, 7.0, 13.0 and 18.0 hours. Besides, the percentage yields were compared in the presence and absence of methanol. This study is useful to determine the optimum reaction time that will produce a maximum yield in hydrolysing the compound especially for ester group.

1.3 Objective

The main objective of this study is:

1. To determine the effect of reaction time on the base hydrolysis of methyl benzoate.
2. To compare the yield of hydrolysis in the presence and absence of methanol.
3. To analyze the hydrolysis product, benzoic acid by using NMR, FTIR and melting point analysis.

2.0 Literature Review

2.1 Ester

Many esters that are known today may either synthetic or natural. According to Ameen and Olatunji (2009), esters have been found in various applications such as in medicines, foods and preservatives. The study stated that one of the important esters; salicylates play a vital role in the food preservatives, pain control and fever control. Besides, acetyl salicylic acid (aspirin) is crucial in preventing the aggregation of platelets and stroke, an important analgesic and anti-inflammatory drugs. Sodium benzoate which is also the derivative of ester have been found to act as food preservative that inhibits microbial growth (Stanojevic *et al.*, 2009). The study done by Ameen and Olatunji (2009) found that the preparations of esters do have some limitation in which it involved lengthy work-ups and the products sometimes contain impurities. Thus, several modifications have been investigated for the preparation of esters in order to reduce the lengthy work-ups and increase the yields of products. One of the methods is by using sulphuric acids as catalyst to remove the water produced during the reaction.

2.1.1 Methyl benzoate

Methyl benzoate is one of the examples of carboxylic acid ester. The general formula for carboxylic acid ester is represented by $R_1C(=O)OR_2$ in which R_1 and R_2 can be alkyl chains, phenyl groups or heteroatoms. The reaction for the acid hydrolysis of carboxylic acid to form ester is the reversible of reaction of ester to form carboxylic acid (Bruice, 2006). It shows that methyl benzoate can be formed by benzoic acid. If ester is the desired product, the reaction should use an excess amount of alcohol in order to drive the reaction reversible. This process is known as Fischer's esterification (Figure 5). According to Francis (2003), methyl benzoate can react with ammonia to form amide. The study found that it is possible to carry

out the reaction in aqueous ammonia due to its higher ability as a nucleophile which is better than water (Figure 6).




Figure 5: Fischer's esterification




Figure 6: Reaction of methyl benzoate with ammonia

2.2 Benzoic acid

According to Bahl and Bahl (2006), benzoic acid was first discovered by Schlee (1775) from gum benzoin. The authors stated that benzoic acid is slightly soluble in cold water but dissolve completely in hot water. The study found that benzoic acid can be prepared in many general methods such as by oxidation of toluene with air in the presence of Co-Mn acetates as catalyst (Figure 7). Researchers also added that benzoic acid can be prepared by passing phthalic anhydride and steam over a metal phthalic catalyst at 200-300°C followed by decarboxylation of benzoic acid from the resulted phthalic acid (Figure 8).




Figure 7: Oxidation of toluene with air




Figure 8: Preparation of benzoic acid from phthalic anhydride

2.1. Application of benzoic acid

Benzoic acid can be used in various applications such as in cosmetics as a pH adjuster and preservative (Nair, 2001). According to Rai *et al.* (2010), benzoic acid a group of food additive important as food preservative and protect the consumers from microbiological risk various bacteria such as *Escherichia coli*, *Listeria monocytogenes*, genus *Aspergillus* and genus *Penicillium* which can cause food poisoning (Rai *et al.*, 2010). The study stated that sodium benzoate which is less toxic is used mostly in acidic food such as salad dressings, carbonated drinks and jams due to its properties which are bacteriostatic and fungistatic under acidic condition of pH below 4.0. According to Bahl and Bahl (2006), benzoic acid plays an important role in medicine as urinary antiseptic and for disinfecting bronchial tubes in vapour form. It also can be used to make aniline blue in dye industry.

3 Hydrolysis

Hydrolysis is a chemical decomposition that involves the breaking of bond and the addition of water (Figure 9) (Ikhazuangbe & Oni, 2015). There are various applications that involved hydrolysis reaction such as hydrolysis of oils and fats to give glycerol and fatty acids (Figure 10) (Salimon *et al.*, 2011). Other organic compound such as carboxylic acid ester, alkyl halide and epoxide also undergo hydrolysis in the medium of acidic, alkaline and neutral (Al-Shamary *et al.*, 2014). Generally, process of hydrolysis is divided into two which are base hydrolysis and acid hydrolysis (Kaye *et al.*, 2004).




Figure 9: Equation of hydrolysis process





Figure 10: Hydrolysis reaction of oils and fats

3.1 Base hydrolysis

Base hydrolysis is an irreversible process. It used base as a catalyst. The common base that used in hydrolysis is NaOH and KOH. KOH based catalyst formulations have been reported to give better yields compared with NaOH based catalyst (Singh *et al.*, 2006). Due to its relationship with soap making, base catalyzed hydrolysis of ester is sometimes known as saponification (Salimon *et al.*, 2011). There are two advantages of doing base hydrolysis. First, it is a one way reaction and next, the product is easier to separate (Salimon *et al.*, 2011). According to a previous study, it has been found that the percentage yield is increase with the increasing time (Salimon *et al.*, 2011). The increasing concentration of catalyst and increase in reaction temperature also increase the productions yield in hydrolysis (Singh *et al.*, 2006).

3.2 Base hydrolysis of carboxylic ester

According to Al-Shamary *et al.* (2014), base hydrolysis of carboxylic ester generally takes place via $\text{S}_{\text{N}}2$ reaction in which the hydroxide ion attacks the carbonyl carbon of an ester to give carboxylic acid and an alcohol (Figure 11). Moreover, the presence of water as solvent and other organic solvent could alter the hydrolytic rate and modify the mechanism of hydrolysis reaction in various ways such as nucleophilic reagent, as a high dielectric constant and as a specific solvating agent for organic reactant and product.



Acid hydrolysis used acid as catalyst. The commonly used acid catalysts are HCl and H_2SO_4 . According to Bruice (2006), when an acid is added to a reaction it protonates the atom in the reactant that has the greatest electron density which is the carbonyl group in ester. Thus, it increases the rate of reaction in tetrahedral intermediate by protonating the carbonyl group. The protonated carbonyl group is more susceptible to nucleophilic attacks compared with non-protonated carbonyl due to the positive charge that is more electrons withdrawing. As a result, it reduces the basicity of the leaving group which makes it easier to eliminate.




Figure 12: Mechanism of acid hydrolysis

3.3 Solvent effects on hydrolysis

As mentioned by Brinker (1988), solvent molecules that formed hydrogen bond to hydroxyl or hydronium ions catalyzed in hydrolysis reduce the catalytic activity under basic

and acidic condition respectively. Thus, the aprotic solvent could make the hydroxyl ions more nucleophilic because it does not form hydrogen bond to hydroxyl ions. In contrast, protic solvent such as alcohol, water and ammonia make hydronium ions more electrophilic. Thus, the presence of MeOH in this study can act as donors and acceptors in hydrogen bonding. The used of organic solvent give several advantages which are increased activity and stability, regiospecificity and stereoselectivity and ease of products recovery (Sharma and Kanwar, 2014).



Figure 1. Conversion of 1-phenylethanol to 1-phenylethyl acetate

3.0 Materials and Methods

3.1 Preparation of 2 M NaOH

80 g of NaOH pellet were weighed by using an electronic balance. The pellets were placed in 500 mL of volumetric flask and distilled water was added to it. After the solution was cooled down, distilled water was added to the volumetric flask again until it reached the mark level.

3.2 Preparation of 2 M KOH

56.11 g of KOH pellet were weighed by using electronic balance. The pellets were placed in 500 mL of volumetric flask and distilled water was added to it. After the solution was cooled down, distilled water was added to the volumetric flask again until it reached the mark level.

3.3 Preparation of 2 M HCl

Distilled water was added to a 200 mL beaker followed by 33 mL of HCl. After the solution cooled down, distilled water was added to the beaker until it reached 200 mL.

3.4 Base hydrolysis of methyl benzoate with solvent




Figure 13: Pathway of hydrolysis of benzoic acid from methyl benzoate

Methyl benzoate (2 mL, 0.02 mol) was added to a 250 mL round bottomed flask and a stirrer bar was placed inside the flask. MeOH (2 mL) was added to the flask and the mixture was stirred until the mixture was homogenous. Next, NaOH/KOH (30mL) was added and the mixture was stirred for 2 hours. After 2 hours, HCl (50 mL) was poured into the mixture